OPTIMAL INSTRUMENT

Difficulty - Baseline

Instructions: Please circle the level of difficulty you have for each activity today.	Able to do without any difficulty	Able to do with little difficulty	Able to do with moderate difficulty	Able to do with much difficulty	Unable to do	Not applicable
1. Lying flat	1	2	3	4	5	9
2. Rolling over	1	2	3	4	5	9
Moving – lying to sitting	1	2	3	4	5	9
4. Sitting	1	2	3	4	5	9
5. Squatting	1	2	3	4	5	9
6. Bending/Stooping	1	2	3	4	5	9
7. Balancing	1	2	3	4	5	9
8. Kneeling	1	2	3	4	5	9
9. Standing	1	2	3	4	5	9
10. Walking-short distance	1	2	3	4	5	9
11. Walking-long distance	1	2	3	4	5	9
12. Walking-outdoors	1	2	3	4	5	9
13. Climbing stairs	1	2	3	4	5	9
14. Hopping	1	2	3	4	5	9
15. Jumping	1	2	3	4	5	9
16. Running	1	2	3	4	5	9
17. Pushing	1	2	3	4	5	9
18. Pulling	1	2	3	4	5	9
19. Reaching	1	2	3	4	5	9
20. Grasping	1	2	3	4	5	9
21. Lifting	1	2	3	4	5	9
22. Carrying	1	2	3	4	5	9


23. From the above list, choose the 3 activities you would most like to be able to do without any difficulty (for example, if you would most like to be able to *climb stairs, kneel*, and *hop* without any difficulty, you would choose: 1.13 2.8 3.14)

1. ____ 2. ____ 3. ____

24. From the above list of 3 activities, choose the primary activity you would most like to be able to do without any difficulty.

Primary goal _____

25. Please mark your pain in the past 24 hours on the scale below:



Copyright © 2012, 2006, 2005 American Physical Therapy Association. All rights reserved.